

Trace Element Contents in Adenocarcinoma of the Human Prostate Gland Investigated by Neutron Activation Analysis

Vladimir Zaichick^{1*} and Sofia Zaichick^{1,2}

¹Radionuclide Diagnostics Department, Medical Radiological Research Centre, Koroleyva St. - 4, Obninsk 249036, Kaluga Region, Russia

²Department of Medicine, College of Medicine, University of Illinois, Chicago, IL 60612, USA

***Corresponding Author:** V. Zaichick, Medical Radiological Research Centre, Koroleyva St. 4, Obninsk 249036, Kaluga Region, Russia; Tel: (48439) 60289; Fax: (495) 956 1440; E-mail: vezai@obninsk.com

Citation: Vladimir Zaichick and Sofia Zaichick (2016) Trace Element Contents in Adenocarcinoma of the Human Prostate Gland Investigated by Neutron Activation Analysis. *Cancer Res Oncol* 1: 002.

Copyright: © 2016 Vladimir Zaichick and Sofia Zaichick. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted Access, usage, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents in normal tissue (n=37) and adenocarcinoma (n=37) of the human prostate gland were investigated using non-destructive instrumental neutron activation analysis with high resolution spectrometry of long-lived radionuclides. Mean values (M ± SEM) for content (mg/kg, dry mass basis) of trace elements in adenocarcinoma were: Ag – 0.242±0.039, Co – 0.0317±0.0032, Cr – 2.31±0.43, Fe – 172±21, Hg – 0.091±0.012, Rb – 9.31±0.88, Sb – 0.445±0.057, Sc – 0.0126±0.0019, Se – 0.629±0.092, and Zn – 127±12. It was observed that in cancerous tissues contents of Co, Rb, Sc, and Zn were significantly lower and those of Ag, Cr, Fe, Hg, and Sb were significantly higher than in normal tissues. The abnormal decrease in levels of Co, Rb, Sc, and Zn in cancerous tissue could be a consequence of malignant transformation. It was supposed that elevated levels of Ag, Cr, Fe, Hg, and Sb as well as an inadequate level of Se are involved in an initiation and a promotion of prostatic adenocarcinoma.

Keywords: Adenocarcinoma of Prostate; Prostate; Trace elements; Neutron activation analysis.

Introduction

Prostate cancer (PCa) is the most common non-cutaneous male cancer in most populations [1]. Although the etiology of PCa is unknown, several risk factors including age and diet (Ca, Zn and some other nutrients) have been well identified [2, 3]. It is also reported that the risk of having PCa drastically increase with age, being three orders of magnitude higher for the age group 40–79 years than for those younger than 39 years [3, 4].

Trace elements have essential physiological functions such as maintenance and regulation of cell function, gene regulation, activation or inhibition of enzymatic reactions, and regulation of membrane function. Essential or toxic (mutagenic, carcinogenic) properties of trace elements depend on tissue-specific need or tolerance, respectively [5]. Excessive accumulation or an imbalance of the trace elements may disturb the cell functions and may result in cellular degeneration or death [5 - 7].

High intraprostatic Zn concentrations are probably one of the main factors acting in both initiation and promotion stages of prostate carcinogenesis [8 - 11]. A significant tendency of age-related increase in Ba, Bi, Cd, Co, Fe, Hg, Pb, Sc, Sn, Th, U, and Zn, mass fraction in the prostate was recently demonstrated by us [12 - 22]. Thus, it seems fair to suppose that besides Zn, many other trace elements also play a role in the pathophysiology of the prostate.

The trace element contents in tissue of the normal [23 - 32] and cancerous [33 - 38] prostate have been studied, producing contradictory results. The majority of these data are based on measurements of processed tissue and in many studies tissue samples are ashed before analysis. In other cases, prostate samples are treated with solvents (distilled water, ethanol etc) and then are dried at a high temperature for many hours. There is evidence that certain quantities of trace elements are lost as a result of such treatment [39 - 41]. Moreover, only a few of these studies employed quality control using certified reference materials for determination of the trace element mass fractions. Thus, the questions about the differences between trace element contents in normal and cancerous prostate tissue remained open.

This work had three aims. The first was to assess the Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, Zn contents in adenocarcinoma of prostate using non-destructive instrumental neutron activation analysis with high resolution spectrometry of long-lived radionuclides (INAA-LLR). The second aim was to measure the levels of trace elements in the nonhyperplastic prostate of healthy men aged over 40 years, and the third was to compare the levels of trace elements in the malignant prostate with those in normal gland.

All studies were approved by the Ethical Committees of the Medical Radiological Research Centre, Obninsk, and the Institute of Forensic Medicine, Moscow.

Materials and Methods

All patients suffered from adenocarcinoma of prostate (n=37, mean age $M \pm SD$ was 63 ± 11 years, range 40-79) were hospitalized in the Urological Department of the Medical Radiological Research Centre. Transrectal puncture biopsy of suspicious indurated regions of the prostate was performed for every patient, to permit morphological study of prostatic tissue at these sites and to estimate their chemical element contents. In all cases the diagnosis adenocarcinoma has been confirmed by clinical and morphological results obtained during studies of biopsy and resected materials.

Normal prostates for the control group samples were removed at necropsy from 37 men (mean age 55 ± 11 years, range 41-87), who had died suddenly. The majority of deaths were due to trauma. A histological examination in the control group was used to control the age norm conformity,

as well as to confirm the absence of microadenomatosis and latent cancer.

All tissue samples were divided into two portions. One was used for morphological study while the other was intended for trace element analysis. After the samples intended for trace element analysis were weighed, they were freeze-dried and homogenized. The sample weighing about 10 mg (for biopsy materials) and 50 mg (for resected materials) was used for chemical element measurement by instrumental neutron activation analysis with high resolution spectrometry of long-lived radionuclides (INAA-LLR). The samples for INAA-LLR were wrapped separately in a high-purity aluminum foil washed with double rectified alcohol beforehand and placed in a nitric acid-washed quartz ampoule.

To determine the contents of the trace elements by comparison with known data for standard, aliquots of commercial, chemically pure compounds, synthetic and natural reference materials were used [42]. Ten sub-samples of international certified reference material (CRM) IAEA H-4 (animal muscle) and ten sub-samples of CRM IAEA HH-1 (human hair) weighing about 50 mg were treated and analyzed under exactly the same conditions as the prostate samples, to allow estimation of the precision and accuracy of results.

The vertical channel of a nuclear reactor was used to determine the samples' content of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn by INAA-LLR. The quartz ampoule containing prostate samples, standards, and certified reference materials was soldered, positioned in an aluminum transport container and exposed to a 24-hour neutron irradiation in the vertical channel with a neutron flux $1.3 \cdot 10^{13} \text{ n} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$. Ten days after the irradiation samples were reweighed and repacked.

Details of the relevant facility for INAA-LLR, nuclear reactions, radionuclides, gamma energies, methods of analysis and the results of quality control were presented in our earlier publications concerning the trace elements of nonhyperplastic human prostate tissue [13].

All prostate samples for INAA-LLR were prepared in duplicate and mean values of trace element contents were used in final calculation. Using the Microsoft Office Excel programs, the summary of statistics, arithmetic mean, standard deviation, standard error of mean, minimum and maximum values, median, percentiles with 0.025 and 0.975 levels was calculated for trace element contents in normal and cancerous prostate tissue. The reliability of difference in the results between the two groups of prostate tissue samples was evaluated by Student's *t*-test. For the estimation of the Pearson correlation coefficient between different pairs of the trace element mass fractions in the normal and cancerous prostate tissue the Microsoft Office Excel program was also used.

Results

Table 1 presents certain statistical parameters (arithmetic mean, standard deviation, standard error of mean, minimal and maximal values, median, percentiles with 0.025 and 0.975 levels) of the Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents in normal prostate tissue and adenocarcinoma of prostate.

The ratios of means and the reliability of difference between mean values of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents in normal and cancerous prostate tissue are presented in Table 2.

Data comparison for mean values of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions in adenocarcinoma tissue of patients of different age group and cancer stage are shown in Table 3 and 4, respectively.

Table 5 depicts results of inter-element correlation calculations (values of r – coefficient of correlation) including all trace elements identified in this work.

The comparison of our results with published data for Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents in normal and cancerous prostate tissue is shown in Table 6.

Table 1: Some statistical parameters of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions (mg/kg, dry mass basis) in normal and cancerous prostate.

Tissue	Parameter	Mean	SD	SEM	Min	Max	Med	Per.0.025	Per.0.975
Normal n=37	Ag	0.0478	0.0456	0.0089	0.0100	0.223	0.0344	0.0100	0.156
	Co	0.0452	0.0237	0.0043	0.0165	0.106	0.0400	0.0169	0.0988
	Cr	0.532	0.422	0.081	0.0300	1.81	0.392	0.0300	1.54
	Fe	111	51	9	35.0	267	100	35.8	221
	Hg	0.056	0.057	0.011	0.0077	0.242	0.0320	0.0114	0.216
	Rb	12.7	5.0	0.9	4.70	25.3	12.0	5.69	24.6
	Sb	0.0445	0.0374	0.0067	0.0046	0.158	0.0380	0.00865	0.154
	Sc	0.0294	0.0236	0.0053	0.00460	0.0771	0.0200	0.00660	0.0768
	Se	0.696	0.240	0.044	0.318	1.43	0.691	0.346	1.15
	Zn	1002	664	109	231	3513	915	252	2185
Adeno- carcinoma n=37	Ag	0.242	0.165	0.039	0.0237	0.527	0.197	0.0293	0.520
	Co	0.0317	0.0152	0.0032	0.00494	0.0674	0.0291	0.0104	0.0623
	Cr	2.31	1.98	0.43	0.165	6.48	1.42	0.255	6.28
	Fe	172	113	21	15.0	427	128	28.5	401
	Hg	0.091	0.055	0.012	0.0159	0.203	0.0797	0.0202	0.201
	Rb	9.31	5.07	0.88	1.00	18.5	8.20	2.12	17.9
	Sb	0.445	0.289	0.057	0.0150	1.03	0.453	0.0355	0.957
	Sc	0.0126	0.0091	0.0019	0.00010	0.0438	0.0107	0.00126	0.0320
	Se	0.629	0.431	0.092	0.0630	1.54	0.507	0.0719	1.43
	Zn	127	74	12	20.0	311	120	20.0	307

M - Arithmetic mean, SD – standard deviation, SEM – standard error of mean, Min – minimum value, Max – maximum value, Med – median, Per.0.025 – percentile with 0.025 level, Per.0.975 – percentile with 0.975 level.

Table 2: Comparison of mean values (M±SEM) of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions (mg/kg, dry mass basis) in normal and cancerous prostate.

Element	Prostatic tissue		Ratio	Student's (t-test)
	Normal	Adenocarcinoma		
	41-87 year	40-79 year	Adenocarcinoma to Normal	$p \leq$
	n=37	n=37		
Ag	0.0478±0.0089	0.242±0.039	5.06	0.00012
Co	0.0452±0.0043	0.0317±0.0032	0.70	0.015
Cr	0.532±0.081	2.31±0.43	4.34	0.00059
Fe	111±9	172±21	1.55	0.012
Hg	0.056±0.011	0.091±0.012	1.63	0.040
Rb	12.7±0.9	9.31±0.88	0.73	0.0076
Sb	0.0445±0.0067	0.445±0.057	10.0	0.0000002
Sc	0.0294±0.0053	0.0126±0.0019	0.43	0.0064
Se	0.696±0.044	0.629±0.092	0.90	0.516 (NS)
Zn	1002±109	127±12	0.13	0.000000016

M - Arithmetic mean, SEM – standard error of mean, NS - not significant difference

Table 3: Differences between mean values (M±SEM) of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions (mg/kg, dry mass basis) in cancerous prostate glands of patients of two age groups.

Element	Adenocarcinoma		Ratio	Student's <i>t</i> -test
	Age group 1	Age group 2		
	40-65 year	66-79 year	Group 2/ Group 1	$p \leq$
	n=18	n=19		
Ag	0.277±0.061	0.213±0.051	0.77	0.438 (NS)
Co	0.0302±0.0042	0.0335±0.0052	1.11	0.635 (NS)
Cr	2.49±0.67	2.10±0.56	0.84	0.660 (NS)
Fe	195±30	156±30	0.80	0.362 (NS)
Hg	0.087±0.013	0.092±0.021	1.06	0.830 (NS)
Rb	10.3±1.2	8.4±1.3	0.82	0.293 (NS)
Sb	0.452±0.065	0.439±0.096	0.97	0.912 (NS)
Sc	0.0147±0.0030	0.0101±0.0021	0.69	0.220 (NS)
Se	0.612±0.099	0.646±0.160	1.06	0.859 (NS)
Zn	131±18	123±17	0.94	0.743 (NS)

M - Arithmetic mean, SEM - standard error of mean, NS - not significant difference.

Table 4: Differences between mean values ($M \pm SEM$) of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions (mg/kg, dry mass basis) in adenocarcinoma of two groups patients with cancer stage in range ($T_1 - T_2$) and ($T_3 - T_4$).

Element	Adenocarcinoma Stage ($T_1 - T_2$) n=12	Adenocarcinoma Stage ($T_3 - T_4$) n=25	Ratio ($T_3 - T_4$)/($T_1 - T_2$)	Student's <i>t</i> -test <i>p</i> ≤
Ag	0.269±0.080	0.228±0.045	0.85	0.666 (NS)
Co	0.0328±0.0091	0.0313±0.0031	0.95	0.877 (NS)
Cr	1.44±0.36	2.74±0.60	1.90	0.079 (NS)
Fe	146±31	185±28	1.27	0.366 (NS)
Hg	0.101±0.021	0.084±0.015	0.83	0.545 (NS)
Rb	11.3±1.7	8.42±0.99	0.75	0.162 (NS)
Sb	0.39±0.10	0.469±0.069	1.20	0.536 (NS)
Sc	0.0107±0.0024	0.0135±0.0026	1.26	0.446 (NS)
Se	0.83±0.20	0.536±0.092	0.65	0.223 (NS)
Zn	179±26	102±10	0.57	0.016

M - Arithmetic mean, SEM - standard error of mean, NS - not significant difference.

Table 5: Intercorrelations (*r* – coefficient of correlation) of pairs of the trace element mass fractions or mass fraction ratios in normal and cancerous prostate glands.

Tissue	Element	Co	Cr	Fe	Hg	Rb	Sb	Sc	Se	Zn
Normal	Ag	-0.15	-0.23	-0.26	-0.06	-0.18	0.28	-0.02	-0.29	0.12
41-87 years n=37	Co	1.00	0.26	0.59 ^b	0.21	0.56 ^b	0.30	0.57 ^b	0.29	0.21
	Cr	0.26	1.00	0.40	0.49 ^a	0.44 ^a	0.18	0.11	0.26	0.08
	Fe	0.59 ^b	0.40	1.00	0.51 ^a	0.66 ^b	0.26	0.70 ^c	0.60 ^b	0.23
	Hg	0.21	0.49 ^a	0.51 ^a	1.00	0.42	0.01	0.62 ^b	0.51 ^a	0.51 ^a
	Rb	0.56 ^b	0.44 ^a	0.66 ^b	0.42	1.00	0.39	0.59 ^b	0.52 ^a	0.28
	Sb	0.30	0.18	0.26	0.01	0.39	1.00	-0.02	-0.03	-0.07
	Sc	0.57 ^b	0.11	0.70 ^c	0.62 ^b	0.59 ^b	-0.02	1.00	0.70 ^c	0.67 ^c
	Se	0.29	0.26	0.60 ^b	0.51 ^a	0.52 ^a	-0.03	0.70 ^c	1.00	0.44 ^a
	Zn	0.21	0.08	0.23	0.51 ^a	0.28	-0.07	0.67 ^c	0.44 ^a	1.00
Adeno- carcinoma 44-87 years	Ag	-0.17	-0.03	-0.01	-0.37	-0.15	0.31	0.36	-0.31	-0.10
	Co	1.00	0.10	0.05	0.05	0.35	-0.43	0.51 ^a	0.39	0.56 ^b
	Cr	0.10	1.00	0.58 ^b	0.28	-0.24	0.16	0.05	-0.12	0.09
	Fe	0.05	0.58 ^b	1.00	0.10	0.14	0.09	0.06	-0.11	-0.10

n=37	Hg	0.05	0.28	0.10	1.00	0.42	-0.20	0.15	0.62 ^b	0.42
	Rb	0.35	-0.24	0.14	0.42	1.00	-0.43	0.26	0.68 ^c	0.27
	Sb	-0.43	0.16	0.09	-0.20	-0.43	1.00	-0.32	-0.42	-0.40
	Sc	0.51 ^a	0.05	0.06	0.15	0.26	-0.32	1.00	0.21	0.18
	Se	0.39	-0.12	-0.11	0.62 ^b	0.68 ^c	-0.42	0.21	1.00	0.78 ^c
	Zn	0.56 ^b	0.09	-0.10	0.42	0.27	-0.40	0.18	0.78 ^c	1.00

Statistically significant difference: ^a - $p \leq 0.05$, ^b - $p \leq 0.01$, ^c - $p \leq 0.001$.

Table 6: Median, minimum and maximum value of means of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fractions (mg/kg, dry mass basis) in normal and cancerous prostate according to data from the literature in comparison with our results (mg/kg, dry mass basis).

Prostate tissue	Element	Published data			This work
		Median of means (n) ^a	Minimum of means M or M±SD, (n) ^b	Maximum of means M or M±SD, (n) ^b	
Normal	Ag	≤0.1 (2)	<0.05 (48) Tipton & Cook 1963	0.2 (7) Tipton et al. 1954	0.048±0.04 6
	Co	0.55 (3)	<0.09 (50) Tipton & Cook 1963	12 (9) Stich 1957	0.045±0.02 4
	Cr	0.56 (3)	0.042 (50) Tipton & Cook 1963	1.4 (8) Tipton et al. 1954	0.53±0.42
	Fe	150 (14)	5.7±0.1 (5) Sangen 1967	1040±65 (10) Jafa et al. 1980	111±51
	Hg	0.65 (1)	0.65±0.58 (5) Liebscher & Smith 1968	0.65±0.58 (5) Liebscher & Smith 1968	0.056±0.05 7
	Rb	34.5(3)	4.7 (9) Stich 1957	58±33 (4) Soman et al. 1970	12.7±5.0
	Sb	0.42 (1)	0.42±0.56 (10) Liebscher & Smith 1968	0.42±0.56 (10) Liebscher & Smith 1968	0.045±0.03 7
	Sc	-	-	-	0.029±0.02 4
	Se	0.625 (7)	0.27 (129) Schöpfer, Drasch & Schrauzer 2010	1.5 (15) Sarafanov et al. 2008	0.70±0.24
	Zn	482 (48)	111 (-) Anspaugh et al. 1973	3218±41 (10) Jafa et al. 1980	1002±664
Adeno-carcinoma	Ag	-	-	-	0.24±0.17
	Co	5.1 (1)	4.7±0.4 (1) Kwiatek et al. 2005	5.4±0.6 (1) Kwiatek et al. 2005	0.032±0.01 5
	Cr	7.0 (4)	0.33±0.06 (1) Kwiatek et al. 2005	217±8 (27) Guntupalli et al. 2007	2.3±2.0
	Fe	195 (9)	12.5±5.0 (23) Kiziler et al. 2010	3530±45 (27) Guntupalli et al. 2007	172±113

Hg	-	-	-	0.091±0.055	
Rb	-	-	-	9.3±5.1	
Sb	-	-	-	0.45±0.29	
Sc	-	-	-	0.013±0.009	
Se	5.5 (7)	0.90±0.55 (29)	Sapota et al. 2009	11.5±3.5 (3) Kwiatek et al. 2004	0.63±0.43
Zn	200 (34)	16.7±3.5 (3)	Kwiatek et al. 2004	840±85 (13) Dhar et al. 1973	127±74

M - Arithmetic mean, SD – standard deviation, (n)^a – number of all references, (n)^b - number of samples.

Discussion

As was shown by us [13] the use of CRM IAEA H-4 and CRM IAEA HH-1 as a certified reference materials for the analysis of samples of prostate tissue can be seen as quite acceptable. Good agreement of the Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents analyzed by INAA-LLR with the certified data of CRMs IAEA indicates an acceptable accuracy of the results obtained in the study of trace elements of the adenocarcinoma of prostate gland presented in Tables 1–6.

The mean values and all selected statistical parameters were calculated for 10 chemical elements (Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn) (Table 1). The mass fraction of these elements were measured in all, or a major portion of normal and cancerous prostate samples.

From Tables 1 and 2, it is observed that in adenocarcinoma the mass fractions of Ag ($p<0.00012$), Cr ($p<0.00059$), Fe ($p<0.012$), Hg ($p<0.040$), and Sb ($p<0.0000002$) are significantly higher while the mass fractions of Co ($p<0.015$), Rb ($p<0.0076$), Sc ($p<0.0064$), and Zn ($p<0.000000016$) are significantly lower than in normal tissues of the prostate. Except for Se, the mass fractions of all trace elements investigated in the study show significant variations in cancerous tissues when compared with normal tissues of the prostate. For example, in adenocarcinoma the Sb mass fractions were 10 times, and the Ag, Cr, Fe, and Hg mass fractions were approximately 1.6 – 5 times, greater than in normal prostate tissue (Table 2). In contrary, the Zn mass fractions were almost 8 times, and the Co, Rb, and Sc mass fractions were approximately 30–60%, lower in adenocarcinoma than in normal prostate tissue (Table 2). It was found that Zn and some other trace elements are involved in functional features of prostate tissue [43]. Because malignant transformation is accompanied by a loss of tissue-specific functional features, this process leads to a significant reduction in the contents of trace elements such as Zn and, probably, Co, Rb, Sc, and Se associated with functional characteristics of the human prostate tissue.

To analyze the effect of age on the trace element mass fractions in the adenocarcinoma of prostate we examined the two age groups; group 1 (samples from patients aged 40 to 65 years) and group 2 (samples from elderly patients aged 66 to 79 years). No age-related differences in the mass fraction of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn in adenocarcinoma were found (Table 3).

To analyze the effect of cancer stage on the trace element mass fractions in the adenocarcinoma of prostate we examined the two group of patients; group 1 (samples from patients with cancer stage ranged T₁ to T₂) and group 2 (samples from patients with cancer stage ranged T₃ to T₄). The mass fraction of Zn in cancer stages ranged T₃–T₄ was almost 2 times lower than in stage T₁ and T₂ (Table 4). Thus, our findings show a positive correlation between the cancer stage and the level of functional losses in malignant transformed prostatic tissue.

In healthy prostate tissue a large proportion (5 out of 9) of statistically significant inter-element correlations were observed for Fe: with Co ($p<0.01$), Hg ($p<0.05$), Rb ($p<0.01$), Sc ($p<0.001$), and Se ($p<0.01$) (Table 5). The positive inter-element correlations of Zn mass fractions with Hg ($p<0.05$), Sc ($p<0.001$), and Se ($p<0.05$) mass fractions were also found in healthy prostate tissue. Additionally, levels of Se have significant relationships with Hg ($p<0.05$), Rb ($p<0.05$), and Sc ($p<0.001$), as well as level of Rb with Co ($p<0.01$), Cr ($p<0.05$), Sc ($p<0.01$), and Se ($p<0.05$). These correlations indicate a possible connection between these elements and the normal function of the prostate. Inter-element correlations between trace elements contents differ markedly in adenocarcinoma of prostate as compared to their relationships in normal prostate tissue. Fe contents have positive correlation only with Cr ($p<0.05$). Zn contents positively correlate with Co ($p<0.01$) and Se ($p<0.001$) whereas Se contents have positive correlation only with Hg ($p<0.01$), Rb ($p<0.001$), and Zn ($p<0.001$). Published data referring to inter-element correlations of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn contents in normal and cancerous tissues of the human prostate gland were not found.

The results for all trace element contents in the prostates of the control group (mean age 55±11 years, range 41-87) are in accordance with our earlier findings in prostates of apparently healthy men aged 41-60 [13]. In data from the literature a number of values for trace element mass fractions in prostate tissue were not expressed on a dry mass basis. Therefore, we calculated these values using published data for water - 80% [44] and ash - 1% on wet mass basis [45] contents in the prostate of adult men. This reported data also includes samples obtained from patients who died from different non-urological diseases. Values obtained for Ag, Cr, Fe, Rb, Se, and Zn contents agree well with median of mean values cited by other researches for the human prostate (Table 6). However, calculated means for Co, Hg, and Sb in this work are an order of magnitude lower than the median of previously reported data. No published data referring to Sc content in the human prostate was found. In the adenocarcinoma of prostate our results were comparable with published data for Cr, Fe and Zn contents and one or two orders of magnitude lower for Co and Se (Table 6). No published data referring to Ag, Hg, Rb, Sb, and Sc contents of cancerous prostate tissue were found.

Compared to other soft tissues, the prostate of young adults has higher levels of Zn and many other trace elements [15 - 18]. Moreover, the level of Zn and some other trace elements continue to increase with age [19 - 22]. In our earlier publications [8, 10, 11] it was discussed in detail that the age-related excessive Zn level in prostatic tissue is probably one of the main factors influencing the initiation and progression of PCa. In addition to the elevated Zn level, an age-related increase and excess in Cr, Fe, Hg, and Sb mass fractions in prostatic tissue may contribute to harmful effects on the gland. There are good reasons for such speculations since many reviews and numerous papers raise the concern about toxicity and carcinogenicity of these metals [46 - 55]. Each of these metals is distinct in its primary mode of action. Moreover, there are several forms of synergistic action of these metals as a part of intracellular metabolism, during which several reactive intermediates and byproducts are created [46, 47, 54]. These reactive species are capable of potent and surprisingly selective activation of

stress-signaling pathways, inhibition of DNA metabolism, repair, and formation of DNA cross links, which are known to contribute to the development of human cancers [47, 48, 52]. In addition to genetic damage via both oxidative and nonoxidative (DNA adducts) mechanisms, metals can also cause significant changes in DNA methylation and histone modifications, leading to alterations in gene expression [48, 49, 51]. In vitro and animal carcinogenic studies provided strong support for the idea that metals can also act as co-carcinogens in combination with nonmetal carcinogens [48].

Our findings show that mass fraction of Ag, Cr, Sb and Zn are significantly different in most adenocarcinomas as compared to normal prostate tissues (Tables 2). Thus, it is plausible to assume that levels of these trace elements in prostate tissue can be used as tumor markers. However, this subjects needs in additional studies.

Conclusion

In this work, trace elemental analysis was carried out in the tissue samples of normal prostate and adenocarcinoma of prostate using INAA-LLR. It was shown that INAA-LLR is an adequate analytical tool for the non-destructive determination of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn content in the tissue samples of human prostate, including needle-biopsy cores. It was observed that in cancerous tissues contents of Co, Rb, Sc, and Zn were significantly lower and those of Ag, Cr, Fe, Hg, and Sb were significantly higher than in normal tissues. In our opinion, the abnormal decrease in levels of Co, Rb, Sc, and Zn in cancerous tissue could be a consequence of malignant transformation. It was supposed that elevated levels of Ag, Cr, Fe, Hg, and Sb as well as an inadequate level of Se are involved in an initiation and a promotion of prostatic adenocarcinoma.

Acknowledgements

We are grateful to Dr Tatyana Sviridova, Medical Radiological Research Center, Obninsk, and Prof. A.A. Zhavoronkov, Institute of Human Morphology, Russian Academy of Medical Sciences, Moscow, for supplying prostate samples.

References

1. Rebbeck TR & Haas GP (2014) 'Temporal trends and racial disparities in global prostate cancer prevalence', *Canadian Journal of Urology*, vol. 21, no. 5, pp. 7496-7506.
2. Yamada K, Araki S, Tamura M, Saka Y, Takahashi M, et al. (2000) 'Epidemiologic determinants of clinically relevant prostate cancer', *International Journal of Cancer*, vol. 89, pp. 259-264.
3. Rebbeck TR (2006) 'Conquering cancer disparities: new opportunities for cancer epidemiology, biomarker, and prevention research', *Cancer Epidemiology, Biomarkers & Prevention*, vol. 15, pp. 1569-1571.
4. Jemal A, Murray T, Samuels A, Ghafoor A, Ward E & Thun MJ (2003) 'Cancer statistics, 2003', *CA: A Cancer Journal For Clinicians*, vol. 53, pp. 5-26.
5. Zaichick V (2006) 'Medical elementology as a new scientific discipline', *Journal of Radioanalytical and Nuclear Chemistry*, vol. 269, pp. 303-309.
6. Ektessabi A, Shikine S, Kitamura N, Rokkum M & Johansson C (2001) 'Distribution and chemical states of iron and chromium released from orthopedic implants into human tissues', *X-Ray Spectrometry*, vol. 30, pp. 44-48.

7. Yoshida S, Ektessabi A & Fujisawa S (2001) 'XAFS spectroscopy of a single neuron from a patient with Parkinson's disease', *Journal of Synchrotron Radiation*, vol. 8, pp. 998–1000.
8. Zaichick V & Zaichick S (1999) 'Role of zinc in prostate cancerogenesis', in *Mengen und Spurenelemente, 19 Arbeitstagung*, eds M Anke et al., Friedrich-Schiller-Universität, Jena, pp.104-115.
9. Leitzmann MF, Stampfer MJ, Wu K, Colditz GA, Willett WC & Giovannucci EL (2003) 'Zinc supplement use and risk of prostate cancer', *Journal of the National Cancer Institute*, vol. 95, pp. 1004-1007.
10. Zaichick V (2004) 'INAA and EDXRF applications in the age dynamics assessment of Zn content and distribution in the normal human prostate', *Journal of Radioanalytical and Nuclear Chemistry*, vol. 262, pp. 229-234.
11. Zaichick V & Zaichick S (2014) 'Age-related histological and zinc content changes in adult nonhyperplastic prostate glands', *Age*, vol. 36, no. 1, pp. 167-181.
12. Zaichick S & Zaichick V (2011a) 'INAA application in the age dynamics assessment of Br, Ca, Cl, K, Mg, Mn, and Na content in the normal human prostate', *Journal of Radioanalytical and Nuclear Chemistry*, vol. 288, no. 1, pp. 197-202.
13. Zaichick S & Zaichick V (2011b) 'The effect of age on Ag, Co, Cr, Fe, Hg, Sb, Sc, Se, and Zn contents in intact human prostate investigated by neutron activation analysis', *Applied Radiation and Isotopes*, vol. 69, no. 6, pp. 827-833.
14. Zaichick V, Nosenko S & Moskvina I (2012) 'The effect of age on 12 chemical element contents in intact prostate of adult men investigated by inductively coupled plasma atomic emission spectrometry', *Biological Trace Element Research*, vol. 147, no. 1-3, pp. 49-58.
15. Zaichick V & Zaichick S (2013a) 'The effect of age on Br, Ca, Cl, K, Mg, Mn, and Na mass fraction in pediatric and young adult prostate glands investigated by neutron activation analysis', *Applied Radiation and Isotopes*, vol. 82, pp. 145-151.
16. Zaichick V & Zaichick S (2013b) 'INAA application in the assessment of Ag, Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn mass fraction in pediatric and young adult prostate glands', *Journal of Radioanalytical and Nuclear Chemistry*, vol. 298, no. 3, pp. 1559-1566.
17. Zaichick V & Zaichick S (2013c) 'NAA-SLR and ICP-AES Application in the assessment of mass fraction of 19 chemical elements in pediatric and young adult prostate glands. *Biological Trace Element Research*, vol. 156, no. 1, pp. 357-366.
18. Zaichick V & Zaichick S (2013d) 'Use of neutron activation analysis and inductively coupled plasma mass spectrometry for the determination of trace elements in pediatric and young adult prostate. *American Journal of Analytical Chemistry*, vol. 4, pp. 696-706.
19. Zaichick V & Zaichick S (2014a) 'INAA application in the assessment of chemical element mass fractions in adult and geriatric prostate glands', *Applied Radiation and Isotopes*, vol. 90, pp. 62-73.
20. Zaichick V & Zaichick S (2014b) 'Determination of trace elements in adults and geriatric prostate combining neutron activation with inductively coupled plasma atomic emission spectrometry', *Open Journal of Biochemistry*, vol. 1, no.2, pp. 16-33.
21. Zaichick V & Zaichick S (2014c) 'Use of INAA and ICP-MS for the assessment of trace element mass fractions in adult and geriatric prostate. *Journal of Radioanalytical and Nuclear Chemistry*, vol. 301, no. 2, pp. 383-397.
22. Zaichick V (2015) 'The variation with age of 67 macro- and microelement contents in nonhyperplastic prostate glands of adult and elderly males investigated by nuclear analytical and related methods', *Biological Trace Element Research*, vol. 168, no. 1, pp. 44-60.
23. Tipton J, Steiner RL, Foland WD, Mueller J & Stanley M (1954) 'USAEC-ORNL-Report-CF-54-12-66'.
24. Stitch SR (1957) 'Trace elements in human tissue. I. A semi-quantitative spectrographic survey', *Biochemistry Journal*, vol. 67, pp. 97-103.
25. Tipton JH & Cook MJ (1963) 'Trace elements in human tissue. Part II. Adult subjects from the United States', *Health Physics*, vol. 9, pp. 103-145.
26. Sangen H (1967) 'The influence of the trace metals upon the aconitase activity in human prostate glands', *The Japanese Journal of Urology*, vol. 58, pp. 1146-1159.
27. Liebscher K & Smith H (1968) 'Essential and nonessential trace elements. A method of determining whether an element is essential or nonessential in human tissue', *Archives of Environmental Health*, vol. 17, pp. 882-891.
28. Soman SD, Joseph KT, Raut SJ, Mulay GD, Parameswaran M & Pandey VK (1970) 'Studies of major and trace element content in human tissues', *Health Physics*, vol. 19, pp. 641-656.
29. Anspaugh LR, Robinson WL, Martin WH & Lowe OA (1973) 'Compilation of Published Information on Elemental Concentrations in human Organs in Both Normal and Diseased States', No. UCRL-51013Pt. 1971-1973, pp. 1-4.
30. Jafa A, Mahendra NM, Chowdhury AR & Kamboj VP (1980) 'Trace elements in prostatic tissue and plasma in prostatic diseases of man', *Indian Journal of Cancer*, vol. 17, pp. 34-37.
31. Sarafanov AG, Todorov TI, Kajdacsy-Balla A, Gray MA, Macias V & Centeno JA (2008) 'Analysis of iron, zinc, selenium and cadmium in paraffin-embedded prostate tissue specimens using inductively coupled plasma mass-spectrometry', *Journal of Trace Elements in Medicine and Biology*, vol. 22, pp. 305-314.
32. Schöpfer J, Drasch G & Schrauzer GN (2010) 'Selenium and cadmium levels and ratios in prostates, livers, and kidneys of nonsmokers and smokers', *Biological Trace Element Research*, vol. 134, pp. 180-187.

33. Kwiatek WM, Banas A, Gajda M, Gałka M, Pawlicki B, Falkenberg G & Cichocki T (2005) 'Cancerous tissues analyzed by SRIXE', *Journal of Alloys and Compounds*, vol. 401, pp. 173-177.
34. Kwiatek WM, Hanson AL, Paluszkiwicz C, Gałka M, Gajda M & Cichocki T (2004) 'Application of SRIXE and XANES to the determination of the oxidation state of iron in prostate tissue sections', *Journal of Alloys and Compounds*, vol. 362, pp. 83-87.
35. Guntupalli JNR, Padala S, Gummuluri AVR, Muktineni RK, Byreddy SR, Sreerama L, Kedarisetti PC, Angalakuduru DP, Satti BR, Venkatathri V, Pullala VBRL & Gavarasana S (2007) 'Trace elemental analysis of normal, benign hypertrophic and cancerous tissues of the prostate gland using the particle-induced X-ray emission technique', *European Journal of Cancer Prevention*, vol. 16, no. 2, pp. 108-115.
36. Kiziler AR, Aydemir B, Guzel S, Alici B, Ataus S, Tuna MB, Durak H & Kilic M (2010) 'May the level and ratio changes of trace elements be utilized in identification of disease progression and grade in prostatic cancer?', *Trace Elements and Electrolytes*, vol. 2, pp. 65-72.
37. Sapota A, Daragó A, Taczalski J & Kilanowicz A (2009) 'Disturbed homeostasis of zinc and other essential elements in the prostate gland dependent on the character of pathological lesions', *Biometals*, vol. 22, pp. 1041-1049.
38. Dhar NK, Goel TC, Dube PC, Chowdhury AR & Kar AB (1973) 'Distribution and concentration of zinc in the subcellular fractions of benign hyperplastic and malignant neoplastic human prostate', *Experimental and Molecular Pathology*, vol. 19, pp. 139-142.
39. Zaichick V (1997) "Sampling, sample storage and preparation of biomaterials for INAA in clinical medicine, occupational and environmental health", in *Harmonization of Health-Related Environmental Measurements Using Nuclear and Isotopic Techniques*, International Atomic Energy Agency, Vienna, pp 123-133.
40. Zaichick V & Zaichick S (1997) 'A search for losses of chemical elements during freeze-drying of biological materials', *Journal of Radioanalytical and Nuclear Chemistry*, vol. 218, no. 2, pp. 249-253.
41. Zaichick, V 2004, 'Losses of chemical elements in biological samples under the dry aching process', *Trace Elements in Medicine*, vol. 5, no. 3, pp. 17-22.
42. Zaichick V (1995) 'Applications of synthetic reference materials in the medical Radiological Research Centre', *Fresenius Journal Analytical Chemistry*, vol. 352, pp. 219-223.
43. Zaichick V & Zaichick S (2014d) 'Androgen-dependent chemical elements of prostate gland', *Andrology & Gynecology: Current Research*, vol. 2, no. 2 *doi:10.4172/2327-4360.1000121* <http://dx.doi.org/10.4172/2327-4360.1000121>
44. Györkey F, Min K-W, Huff JA & Györkey P (1967) 'Zinc and magnesium in human prostate gland: Normal, hyperplastic, and neoplastic', *Cancer Research*, vol. 27, pp. 1349-1353.
45. Saltzman BE, Gross SB, Yeager DW, Meiners BG & Gartside PS (1990) 'Total body burdens and tissue concentrations of lead, cadmium, copper, zinc, and ach in 55 human cadavers', *Environmental Research*, vol. 52, pp. 126-145.
46. Sunderman FW (1979) 'Mechanism of metal carcinogenesis', *Biological Trace Element Research*, vol. 1, pp. 63-86.
47. Snow ET (1992) 'Metal carcinogenesis: mechanistic implications', *Pharmacology & Therapeutics*, vol. 53, pp. 31-65.
48. Salnikow K & Zhitkovich A (2008) 'Genetic and epigenetic mechanisms in metal carcinogenesis and cocarcinogenesis: nickel, arsenic, and chromium', *Chemical Research in Toxicology*, vol. 21, pp. 28-44.
49. Toyokuni S (2009) 'Role of iron in carcinogenesis: cancer as a ferrotoxic disease', *Cancer Science*, vol. 100, pp. 9-16.
50. Martinez-Zamudio R & Ha HC (2011) 'Environmental epigenetics in metal exposure', *Epigenetics*, vol. 6, pp. 820-827.
51. Tokar EJ, Benbrahim-Tallaa L & Waalkes MP (2011) 'Metal ions in human cancer development', *Metal Ions in Life Sciences*, vol. 8, pp. 375-401.
52. Chervona Y, Arita A & Costa M (2012) 'Carcinogenic metals and the epigenome: understanding the effect of nickel, arsenic, and chromium', *Metallomics*, vol. 4, pp. 619-627.
53. Tchounwou PB, Yedjou CG, Patlolla AK & Sutton DJ (2012) 'Heavy metal toxicity and the environment', *Molecular, Clinical and Environmental Toxicology*, vol. 101, pp. 133-164.
54. Koedrith P, Kim H, Weon JI & Seo YR (2013) 'Toxicogenomic approaches for understanding molecular mechanisms of heavy metal mutagenicity and carcinogenicity', *International Journal of Hygiene and Environmental Health*, vol. 216, pp. 587-598.
55. Tabrez S, Priyadarshini M, Priyamvada S, Khan MS, Na A & Zaidi SK (2014) 'Gene-environment interactions in heavy metal and pesticide carcinogenesis', *Mutation Research*, vol. 760, pp. 1-9.

Please Submit your Manuscript to Cresco Online Publishing

<http://crescopublications.org/submitmanuscript.php>